

STABILITY OF RELATIVISTIC FORCE-FREE JETS

RAMESH NARAYAN¹, JASON LI² AND ALEXANDER TCHEKHOVSKOY³*Draft version January 30, 2009*

ABSTRACT

We consider a two-parameter family of cylindrical force-free equilibria, modeled to match numerical simulations of relativistic force-free jets. We study the linear stability of these equilibria, assuming a rigid impenetrable wall at the outer cylindrical radius R_j . We find that equilibria in which the Lorentz factor $\gamma(R)$ increases monotonically with increasing radius R are stable. On the other hand, equilibria in which $\gamma(R)$ reaches a maximum value at an intermediate radius and then declines to a smaller value γ_j at R_j are unstable. The most rapidly growing mode is an $m = 1$ kink instability which has a growth rate $\sim (0.4/\gamma_j)(c/R_j)$. The e -folding length of the equivalent convected instability is $\sim 2.5\gamma_j R_j$. For a typical jet with an opening angle $\theta_j \sim \text{few}/\gamma_j$, the mode amplitude grows weakly with increasing distance from the base of the jet, much slower than one might expect from a naive application of the Kruskal-Shafranov stability criterion.

Subject headings: instabilities – MHD – galaxies: jets

1. INTRODUCTION

Relativistic jets in astrophysical sources have been known for many decades. Although their enormous power, large Lorentz factor and strong collimation have been widely studied, these phenomena still lack an accepted explanation. An even greater mystery is the remarkable coherence and apparent stability of jets over very large length scales. This is the topic of the present paper.

The most promising models of relativistic jets involve acceleration and collimation by magnetic fields with footpoints attached to a spinning black hole or neutron star or accretion disk. The forced rotation of the field lines induces a strong toroidal component of the magnetic field, which is responsible for accelerating the jet (e.g., Narayan, McKinney, & Farmer 2007; Tchekhovskoy, McKinney, & Narayan 2008, hereafter TMN08; and references therein). In this picture the toroidal component of the field dominates over other field components.

According to the well-known Kruskal-Shafranov criterion (e.g., Bateman 1978), cylindrical magnetohydrodynamic (MHD) configurations in which the toroidal field dominates are violently unstable to the $m = 1$ kink instability (also called the screw instability). The KS criterion for instability is

$$\left| \frac{B_\phi}{B_p} \right| > \frac{2\pi R_j}{z}, \quad (1)$$

where B_ϕ and B_p are the toroidal and poloidal magnetic field strengths, R_j is the cylindrical radius of the jet and z is the length of the jet (i.e., distance from the base of the jet). Typical jet models, including the ones described in this paper (see §2.2), have $B_\phi \sim \gamma_j B_p$, where $\gamma_j \gg 1$ is the Lorentz factor of the jet, and they have jet angles $\theta_j \sim R_j/z \sim \text{few}/\gamma_j$. Substituting these scalings in equation (1), it is obvious that the KS instability criterion is easily satisfied in relativistic jets. We therefore expect astrophysical jets to be violently unstable, as argued for example by Begelman (1998) and Li (2000). Yet, jets in nature are apparently quite stable. How is this possible?

Many authors have investigated this question. They have used jet models with a variety of velocity profiles, geometrical shapes, composition and boundary conditions (Kadomtsev 1966; Bateman 1978; Ferrari et al. 1978; Benford 1981; Payne & Cohn 1985), and applied both analytical and numerical methods (Istomin & Pariev 1996; Begelman 1998; Lyubarskii 1999; Li 2000; Lery et al. 2000; Appl et al. 2000; Tomimatsu et al. 2001; Mizuno et al. 2007; Moll et al. 2008; McKinney & Blandford 2008). As a result of this large body of work, several kinds of unstable modes have been identified: reflection modes, Kelvin-Helmholtz modes, current-carrying modes, etc. Unfortunately, it is difficult to synthesize the results and extract universal principles.

A fruitful approach in this field is to reduce relativistic jet models to their barest minimum. One such approach is to consider *force-free* jet models in which one ignores the inertia and pressure of the plasma and considers only charges, currents and fields. The force-free approximation is valid whenever the energy density in fields dominates over matter energy density, as in pulsar magnetospheres (Goldreich & Julian 1969; Ruderman & Sutherland 1975). The force-free approximation is valid also in relativistic MHD jets, at least inside the fast surface (e.g., Tchekhovskoy et al. 2009).

Theoretical studies of force-free jets have led to the identification of two distinct stability criteria. In a detailed analysis, Istomin & Pariev (1996) showed that cylindrical force-free jets in which B_z is independent of R are stable. Lyubarskii (1999) considered models with non-constant B_z and showed that force-free jets are unstable if

$$\frac{dB_z}{dR} < 0, \quad (2)$$

i.e., if the poloidal field decreases with increasing distance from the axis. We refer to equation (2) as the IPL criterion for instability. On the other hand, Tomimatsu, Matsuoka, & Takahashi (2001) showed via an approximate analysis⁴ that force-free jets are unstable if

$$\left| \frac{B_\phi}{B_p} \right| > \frac{\Omega R}{c}, \quad (3)$$

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, rnarayan@cfa.harvard.edu

² Department of Astrophysical Sciences, Princeton University, Peyton Hall, Ivy Lane, Princeton, NJ 08544, jgli@princeton.edu

³ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, atchekho@cfa.harvard.edu

⁴ They effectively restricted their analysis to the region of the jet inside the light cylinder. Therefore, the flow speeds they considered were only quasi-relativistic at best.

where Ω is the angular velocity of the field line. This criterion — the TMT criterion — differs from the KS criterion (1) in that it explicitly accounts for rotation. It is also apparently very different from the IPL criterion.

We describe in this paper a class of force-free cylindrical jet equilibria which closely match the numerical force-free jet simulations reported in TMN08. Within the context of force-free jets from rigidly-rotating stars, we believe that this two-parameter family of equilibria is generic and fairly complete. We study the stability properties of these equilibria and attempt to relate our results to the KS, IPL and TMT criteria (eqs. 1, 2 and 3).

In §2 we summarize the numerical simulation results of TMN08 (§2.1) and we describe an analytical force-free jet model which matches the simulation data very closely (§2.2). In §3 we carry out a linear stability analysis of these equilibria and show that the linear modes of the system are obtained by solving an eigenvalue problem involving two coupled differential equations, with appropriate boundary conditions. In §4 we numerically solve the equations and identify the unstable modes in the system. We also derive an approximate estimate for the growth rate of the instability. We conclude in §5 with a summary and discussion. We use (r, θ, ϕ) for spherical coordinates and (R, ϕ, z) for cylindrical coordinates.

2. FORCE-FREE JET EQUILIBRIUM

2.1. Structure of Force-Free Jets

TMN08 considered a rigidly-rotating star of unit radius ($r = 1$) surrounded by a differentially-rotating infinitely thin disk extending from $R = 1$ outward. The star was threaded by a uniform radial magnetic field B_r , and the disk was threaded by a power-law distribution of vertical field,

$$B_z \propto R^{\nu-2}. \quad (4)$$

Using a relativistic force-free code (Gammie et al. 2003; McKinney & Gammie 2004; McKinney 2006; Mignone & McKinney 2007; Tchekhovskoy et al. 2007), TMN08 numerically evolved the system and obtained the equilibrium configuration of the magnetic field.

Following TMN08, we will call the field lines that emerge from the star as the “jet” and the field lines from the disk as the “wind.” The critical field line that emerges from the star-disk boundary defines the boundary between the jet and the wind. This boundary starts off at $\theta = \pi/2$ at the surface of the star ($r = R = 1$) but decreases to smaller values of θ with increasing r (or z).

We are primarily interested in the “jet” — the bundle of field lines attached to the central star. Since all of these field lines rotate at the angular velocity Ω of the star, the Alfvén surface for these lines takes the form of a cylinder — the “light cylinder” — with radius $R_A = c/\Omega$. Field lines become strongly toroidal once they are outside the Alfvén surface, which is where most of the collimation and acceleration occurs (TMN08).

TMN08 showed that the structure of the jet is strongly affected by the radial pressure profile of the region surrounding the jet. Specifically, if we write the radial variation of the confining pressure as $p \propto r^{-\alpha}$, the jet properties are determined by the value of α . In the numerical experiments, the pressure was caused by a force-free disk wind, and α was determined by the index ν defined in equation (4) according to

$$\alpha = 2(2 - \nu). \quad (5)$$

At distance z along the axis, the cylindrical radius R_j of the jet is approximately given by

$$R_j \sim z^{\alpha/4}. \quad (6)$$

For all $\alpha < 4$, the jet collimates as it moves away from the star (Lynden-Bell 2006). As a result, at a sufficiently large distance from the central star ($r \gg 1$), the jet is nearly cylindrical in shape.

In the asymptotic nearly-cylindrical region of the jet, force balance in the R direction is described by the following equation (TMN08):

$$\frac{d}{dR} \left(\frac{B^2 - E^2}{8\pi} \right) + \left(\frac{B_\phi^2 - E^2}{4\pi R} \right) + \left(\frac{B_p^2 - E^2}{4\pi R_c} \right) = 0, \quad (7)$$

where B is the total magnetic field strength, B_ϕ is the toroidal field strength, and B_p is the poloidal field strength:

$$B = \sqrt{B_p^2 + B_\phi^2}, \quad B_p = \sqrt{B_R^2 + B_z^2}. \quad (8)$$

The electric field $\vec{E}$ is given by

$$\vec{E} = -(\Omega R/c) \hat{\phi} \times \vec{B}, \quad (9)$$

where Ω is the angular velocity of the field line. The electric field has only a poloidal component: $E_p = \Omega R B_p / c$.

Each of the three terms on the left-hand side of equation (7) represents a force in the $-R$ direction. The quantity $(B^2 - E^2)/8\pi$ is the pressure of the force-free fluid in the moving frame; therefore, the first term describes the inward force due to the gradient of pressure. The second term arises from the toroidal curvature of the field line. The toroidal magnetic field B_ϕ contributes an inward force due to “hoop stress,” while the poloidal electric field E contributes an outward force.⁵ The third term in (7) gives analogous contributions from the poloidal curvature of the field line, where R_c is the poloidal radius of curvature; once again there is an inward force due to the poloidal magnetic hoop stress and an outward force due to the electric field. Note that the contributions involving E are important only for relativistic flows. In standard non-relativistic MHD, one neglects these terms and keeps only the terms involving B .

TMN08 showed that models with $\alpha > 2$, i.e., $\nu < 1$, are good analogs of relativistic jets found in nature (especially the jets of gamma-ray burst). Figure 2.1 shows numerical results corresponding to the “fiducial” force-free simulation in TMN08 with $\nu = 0.75$ (equivalent to $\alpha = 2.5$). Panels (a) and (b) show results for a relatively near region of the jet at $z = 10^2$, and panels (c) and (d) show results for a more distant region at $z = 10^7$. In each case, the abscissa corresponds to the cylindrical radius R normalized by the local “jet radius” R_j , which is the cylindrical radius of the last jet field line that separates the jet from the surrounding disk wind.

The main features of the numerical solution are as follows. First, from panels (a) and (c) we see that B_p is essentially constant inside the jet, showing hardly any variation with R . As we show in Appendix A, this is required for force-free jet solutions that smoothly connect to the central compact object.⁶

⁵ By equation (9), $\vec{E}$ is directly proportional to Ω , so the latter term results from rotation and may loosely be viewed as a sort of “centrifugal force” (V. Beskin, private communication).

⁶ Asymptotic force-free jet configurations with non-constant profiles of B_p are certainly possible (Istomin & Pariev 1996), but there exists no solution that would smoothly connect them to the compact object.




FIG. 1.— Numerical results from a force-free jet simulation with $\alpha = 2.5$ (“fiducial model” corresponding to $\nu = 0.75$ in TMN08). [Panel (a)]: Field components as functions of normalized cylindrical radius R/R_j at $z = 10^2$. The solid horizontal line shows $B_p^2(R)/B_p^2(0)$, the short-dashed line shows $[B_\phi^2(R) - E^2(R)]/B_p^2(0)$ and the long-dashed line shows $E^2(R)/B_p^2(0)$. The dotted lines are the corresponding results for the analytical model with $\gamma_m = 6$, $R_m = 1.2$ (Model A, §2.2.2). The vertical solid line shows the boundary between the jet and the external confining medium. [Panel (b)]: Lorentz factor $\gamma(R)$ (solid line), and the two approximations, $\gamma_1(R)$ (short-dashed line) and $\gamma_2(R)$ (long-dashed line), at the same z . The dotted lines are from the analytical model. [Panels (c), (d)]: Similar to (a), (b), but at $z = 10^7$. The dotted lines in these panels correspond to the analytical model with $\gamma_m = 2600$, $R_m = 0.18$ (Model C, §2.2.2).

Second, since Ω is constant and $E \propto \Omega R B_p$ (eq. 9), we have $E^2 \propto R^2$. Third, B_ϕ is almost equal to E and so B_ϕ also varies primarily as R^2 . Fourth, $|B_\phi|$ is slightly larger than E with $B_\phi^2 - E^2 \propto R^4$. Because of this property, the first two terms in equation (7) both give an inward force. Therefore, we obtain the fifth feature of the solution, viz., the third term in (7), which involves the poloidal curvature of the field line, is important for force balance. This is the only outward force in the balance equation – it is outward because E is of order B_ϕ and is much greater than B_p outside the light cylinder. This force has to balance the other two terms.

The velocity of a force-free flow is usually identified with the drift velocity,

$$\frac{\vec{v}}{c} = \frac{\vec{E} \times \vec{B}}{B^2}, \quad (10)$$

and the Lorentz factor is defined correspondingly. Since $\vec{E} \cdot \vec{B} = 0$, it is easily shown that

$$\gamma^2 = \frac{B^2}{B^2 - E^2}. \quad (11)$$

Panels (b) and (d) show the variation of γ as a function of the normalized cylindrical radius R/R_j in the numerical model.

TMN08 derived two approximate relations for γ ,

$$\gamma_1 = [1 + (\Omega R/c)^2]^{1/2}, \quad (12)$$

$$\gamma_2 = (3R_c/R)^{1/2}, \quad (13)$$

and they showed that the net γ of the fluid is given to good accuracy by the following simple formula,

$$\frac{1}{\gamma^2} = \frac{1}{\gamma_1^2} + \frac{1}{\gamma_2^2}. \quad (14)$$

Along each field line, γ is initially determined mainly by rotation, and so $\gamma \approx \gamma_1$. This is a region of efficient acceleration which TMN08 called the *first acceleration regime*. However, beyond a certain distance from the star, the effect of poloidal field line curvature becomes important, and γ switches to the less efficient γ_2 , the *second acceleration regime*.

Relatively near the star, all field lines in the jet are in the first acceleration regime and $\gamma(R)$ behaves like γ_1 (eq. 12), as seen in panel (b). Specifically, γ increases more or less linearly with R and reaches its maximum value at the edge of the jet at $R = R_j$. However, when we consider the jet at a larger distance from the star, some of the field lines have already switched to the second acceleration regime, where $\gamma \sim \gamma_2 \propto 1/R$ (see eq. 13, coupled with eq. 21 below). We then have the results

shown in panel (d), where the maximum Lorentz factor occurs at some radius R_m inside the jet, not at the boundary; we have $\gamma \sim \gamma_1 \propto R$ for $R \lesssim R_m$ and $\gamma \sim \gamma_2 \propto R^{-1}$ for $R_m \lesssim R < R_j$. TMN08 discuss in detail the physics of the two acceleration regimes.

2.2. Analytical Jet Model

The axisymmetric numerical jets models described in §2.1 have magnetic and electric field components that are functions of both R and z . This is not convenient for linear perturbation analysis. Since the numerical models are nearly cylindrical at large distance (i.e., $dR/dz \ll 1$), we consider now an idealized jet equilibrium model which is perfectly cylindrical and in which all quantities are functions only of R . We choose the following specific functional forms:

$$B_{0R} = 0, \quad (15)$$

$$B_{0\phi} = -[2(\gamma_m^2 - 1)(R/R_m)^2 + (R/R_m)^4]^{1/2} \equiv -f(R), \quad (16)$$

$$B_{0z} = \exp[-3R^2/4(\gamma_m^2 - 1)R_m^2] \equiv g(R), \quad (17)$$

$$E_{0R} = -[2(\gamma_m^2 - 1)]^{1/2}(R/R_m) \equiv -h(R), \quad (18)$$

$$E_{0\phi} = 0, \quad (19)$$

$$E_{0z} = 0, \quad (20)$$

$$R_c = 2(\gamma_m^2 - 1)R_m^2/3R. \quad (21)$$

The zeros in the subscripts are meant to indicate that all these quantities refer to the unperturbed model. The model has two parameters, γ_m and R_m , whose meanings are explained below. For simplicity, we have chosen units such that $B_{0z} = 1$ at the jet axis ($R = 0$). Also, we have assumed that B_{0z} and Ω are positive, so both $B_{0\phi}$ and E_{0R} are negative, i.e., magnetic field lines are swept backward with respect to the rotation and the electric field is pointed radially inward. With this choice of signs, the three functions, $f(R)$, $g(R)$ and $h(R)$, are positive. Note that, in all cases of interest, $g(R)$ is practically equal to unity. The particular exponential form given in equation (17) is designed to handle small higher-order terms in the force balance equation (22), but the deviations of $g(R)$ from unity are tiny and unimportant.⁷

By direct substitution it is easily verified that the above model satisfies the radial force balance equation (7). Under cylindrical symmetry, this equation takes the form:

$$\frac{d}{dR} \left(\frac{B_{0\phi}^2 + B_{0z}^2 - E_{0R}^2}{8\pi} \right) + \left(\frac{B_{0\phi}^2 - E_{0R}^2}{4\pi R} \right) + \left(\frac{B_{0z}^2 - E_{0R}^2}{4\pi R_c} \right) = 0. \quad (22)$$

The first two terms are positive, i.e., both represent inward forces, with the first term providing twice as much force as the second. The third term is negative and its magnitude is equal to the sum of the other two terms.

An important feature of the above model is that the outward force from the third term involves the poloidal curvature radius R_c . Technically, a perfectly cylindrical model has $R_c \rightarrow \infty$. To get around this problem, we treat R_c as an externally imposed property of the solution which is adjusted so as to reproduce the poloidal curvature force present in the numerical jet model. In other words, even though we have straightened out field lines in the z -direction by enforcing cylindrical geometry, we still retain the effect of poloidal curvature by means of an artificial external force. This procedure

⁷ As a test, in the stability analysis described later we have done the calculations both with the full expression for $g(R)$ given in eq. (17) and with the simpler choice $g(R) = 1$. The results are practically the same.

is analogous to the widely-used shearing sheet approximation in accretion disk studies (Goldreich & Tremaine 1978; Narayan et al. 1987) in which fluid streamlines are straightened out in the azimuthal direction, but the effect of azimuthal curvature is still retained via a Coriolis force. Note that, apart from the extra term due to poloidal field curvature, equation (7) is identical to the standard balance condition derived in other papers in the literature, e.g., equation (6) in Istomin & Pariev (1996) or equation (16) in Lyubarskii (1999).

To get a better idea of the nature of the above analytical model, we now make a couple of simplifications. As already mentioned, B_{0z} is practically independent of R inside the jet. Also, for highly relativistic jets, we invariably have $B_{0\phi}^2 - E^2 \ll B_{0\phi}^2$. We therefore replace equations (16), (17) and (18) by the following simpler formulae

$$B_{0z} \approx 1, \quad (23)$$

$$B_{0\phi} \approx E_{0R} = -[2(\gamma_m^2 - 1)]^{1/2}(R/R_m), \quad (24)$$

$$B_{0\phi}^2 - E^2 = (R/R_m)^4. \quad (25)$$

By equation (9), the angular velocity of rotation of the field lines is given by

$$\Omega = -cE_{0R}/B_{0z}R \approx \sqrt{2(\gamma_m^2 - 1)}(c/R_m), \quad (26)$$

and is the same for all field lines, as required for a rigidly rotating star at the base of the jet.⁸ Using these simpler expressions, we obtain the following result for the Lorentz factor:

$$\begin{aligned} \frac{1}{\gamma^2(R)} &= \frac{B_0^2 - E_0^2}{B_0^2} = \frac{B_{0\phi}^2 + B_{0z}^2 - E_{0R}^2}{B_{0\phi}^2 + B_{0z}^2} \\ &\approx \frac{1 + (R/R_m)^4}{1 + 2(\gamma_m^2 - 1)(R/R_m)^2} \end{aligned} \quad (27)$$

$$\approx \frac{1}{1 + (\Omega R/c)^2} + \frac{R}{3R_c}, \quad (28)$$

where we have made use of equations (21) and (26). We thus reproduce the result given earlier in equation (14).

It is easily shown that γ reaches a maximum at $R = R_m$ and that its value at this radius is equal to γ_m . Thus, the two model parameters R_m and γ_m allow us to control the basic features of the equilibrium.

2.2.1. $R_m > R_j$: Maximum Lorentz Factor Located at the Jet Boundary

A jet in which all field lines are in the first acceleration regime has its maximum Lorentz factor at the boundary of the jet, $R = R_j$. This corresponds to choosing $R_m > R_j$ in the analytical model, so that the term $(R/R_m)^4$ in the numerator of equation (27) can be neglected. In this case, the profile of γ has two segments: the region of the jet inside the light cylinder ($R < R_A = c/\Omega$) which is not accelerated very much, and the region outside the light cylinder which has γ increasing linearly with radius,

$$\gamma(R) \approx \begin{cases} 1, & R < R_A = R_m/\sqrt{2(\gamma_m^2 - 1)}, \\ R/R_A \approx (R/R_j)\gamma_j, & R_A < R < R_j, \end{cases}$$

⁸ Since we have designed our analytical model to match the numerical models of TMN08, all of our models have constant $\Omega(R)$. It would be straightforward to generalize the model to non-constant $\Omega(R)$, using additional parameters.




FIG. 2.— Profiles of γ vs R/R_j for the analytic Models A, B and C.

where γ_j is the Lorentz factor at the jet boundary,

$$\gamma_j \approx \gamma_1(R_j) = [1 + 2(\gamma_m^2 - 1)(R_j/R_m)^2]^{1/2}. \quad (29)$$

The dotted lines in panels (a) and (b) in Fig. 2.1 show the dependences of various quantities as a function of R/R_j for a model with $\gamma_m = 6$ and $R_m = 1.2R_j$. The agreement with the numerical simulation results at $z = 10^2$ is striking. We call the analytical model with this particular choice of γ_m and R_m as Model A:

$$\text{Model A:} \quad \gamma_m = 6, \quad R_m = 1.2R_j. \quad (30)$$

The solid line in Fig. 2 shows the variation of γ as a function of R for this model.

2.2.2. $R_m < R_j$: Maximum Lorentz Factor Located Inside the Jet

As we described in §2.1, a jet in which some field lines have switched to the second acceleration regime has its maximum Lorentz factor inside the jet. This means $R_m < R_j$. In this case, the Lorentz factor γ_j at the jet boundary is roughly equal to

$$\gamma_j \approx \gamma_2(R_j) = [2(\gamma_m^2 - 1)]^{1/2} R_m/R_j. \quad (31)$$

Now the profile of γ has three segments:

$$\gamma(R) \approx \begin{cases} 1, & R < R_A, \\ R/R_A \approx (R/R_m)\gamma_m, & R_A < R < R_m, \\ (R_m/R)\gamma_m \approx (R_j/R)\gamma_j, & R_m < R < R_j. \end{cases} \quad (32)$$

The dotted lines in panels (c) and (d) of Fig. 2.1 show model results corresponding to $\gamma_m = 2600$ and $R_m = 0.18R_j$. We find excellent agreement with the numerical results at $z = 10^7$. We call the analytical model with these values of γ_m and R_m as Model C. For completeness we also consider a less extreme model called Model B in which $\gamma_m = 6$ and $R_m = 0.3R_j$:

$$\text{Model B:} \quad \gamma_m = 6, \quad R_m = 0.3R_j, \quad (33)$$

$$\text{Model C:} \quad \gamma_m = 2600, \quad R_m = 0.18R_j. \quad (34)$$

The dashed and dotted lines in Fig. 2 show the variations of γ as a function of R for these two models.

3. LINEAR PERTURBATION ANALYSIS

We now consider linear perturbations of the cylindrical equilibrium described in §2.2. The unperturbed state has magnetic and electric fields

$$\vec{B}_0 = B_{0R}\hat{R} + B_{0\phi}\hat{\phi} + B_{0z}\hat{z}, \quad (35)$$

$$\vec{E}_0 = E_{0R}\hat{R} + E_{0\phi}\hat{\phi} + E_{0z}\hat{z}, \quad (36)$$

where the various components are given by the expressions in equations (15)–(20). As mentioned previously, we choose B_{0z} and Ω to be positive, so $B_{0\phi}$ and E_{0R} are negative. The unperturbed current and electric charge are

$$\vec{J}_0 = \frac{c}{4\pi} \vec{\nabla} \times \vec{B}_0 = \frac{c}{4\pi} \left[- \left(\frac{dB_{0z}}{dR} + \frac{B_{0z}}{R_c} \right) \hat{\phi} + \frac{1}{R} \frac{d}{dR} (RB_{0\phi}) \hat{z} \right],$$

$$\rho_0 = \frac{1}{4\pi} \vec{\nabla} \cdot \vec{E}_0 = \frac{1}{4\pi R} \frac{d}{dR} (RE_{0R}) + \frac{1}{4\pi} \frac{E_{0R}}{R_c}.$$

Note that we have included terms involving R_c in the unperturbed current and charge density. These terms describe the contributions of poloidal field curvature to the quantities $\vec{\nabla} \times \vec{B}_0$ and $\vec{\nabla} \cdot \vec{E}_0$, respectively. By including these terms, we retain the forces associated with poloidal field curvature without actually having curved field lines in the model.

3.1. The Eigenvalue Problem

We now consider small perturbations. Let us write the perturbed electric field as $\vec{E} = \vec{E}_0 + \vec{E}_1$, where $\vec{E}_1$ is a small perturbation of the form

$$\vec{E}_1 = [E_{1R}(R)\hat{R} + E_{1\phi}(R)\hat{\phi} + E_{1z}(R)\hat{z}] \exp(-i\omega t + im\phi + ikz). \quad (37)$$

Let us similarly write $\vec{B} = \vec{B}_0 + \vec{B}_1$, $\vec{J} = \vec{J}_0 + \vec{J}_1$, $\rho = \rho_0 + \rho_1$. Each of these small perturbations can be expressed in terms of the perturbed electric field via Maxwell's equations. From

$$\frac{1}{c} \frac{\partial \vec{B}}{\partial t} = -\vec{\nabla} \times \vec{E}, \quad (38)$$

we obtain

$$\vec{B}_1 = -\frac{ic}{\omega} \vec{\nabla} \times \vec{E}_1. \quad (39)$$

From

$$\frac{1}{c} \frac{\partial \vec{E}}{\partial t} = \vec{\nabla} \times \vec{B} - \frac{4\pi}{c} \vec{J}, \quad (40)$$

we obtain

$$\vec{J}_1 = \frac{i\omega}{4\pi} \vec{E}_1 - \frac{ic^2}{4\pi\omega} \vec{\nabla} \times (\vec{\nabla} \times \vec{E}_1). \quad (41)$$

Finally, from

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho, \quad (42)$$

we obtain

$$\rho_1 = \frac{1}{4\pi} \vec{\nabla} \cdot \vec{E}_1. \quad (43)$$

Since the perturbed system is force-free, it must satisfy $\vec{E} \cdot \vec{B} = 0$. The zeroth order terms satisfy this trivially (as they should). The first order terms give the condition $\vec{E}_1 \cdot \vec{B}_0 + \vec{E}_0 \cdot \vec{B}_1 = 0$. Substituting for the various quantities, this condition allows us to solve for $E_{1\phi}$ in terms of E_{1z} :

$$E_{1\phi} = C_1 E_{1z}, \quad (44)$$

where the function C_1 is given by

$$C_1 = \frac{\omega R g - c m h}{\omega R f - c k R h}. \quad (45)$$

Successive differentiations give

$$E'_{1\phi} = C_1 E'_{1z} + C'_1 E_{1z}, \quad (46)$$

$$E''_{1\phi} = C_1 E''_{1z} + 2C'_1 E'_{1z} + C''_1 E_{1z}. \quad (47)$$

We now consider the force balance condition: $\rho \vec{E} + (1/c) \vec{J} \times \vec{B} = 0$. The zeroth order terms give

$$\rho_0 \vec{E}_0 + \frac{1}{c} \vec{J}_0 \times \vec{B}_0 = 0, \quad (48)$$

which is simply the equilibrium force balance condition (22). Notice that the poloidal curvature terms in $\vec{J}_0$ and ρ_0 are necessary to satisfy equilibrium in the unperturbed solution. From the first order terms in the force balance equation we obtain

$$\rho_1 \vec{E}_0 + \rho_0 \vec{E}_1 + \frac{1}{c} \vec{J}_1 \times \vec{B}_0 + \frac{1}{c} \vec{J}_0 \times \vec{B}_1 = 0. \quad (49)$$

The $\hat{\phi}$ component of this equation gives a relation between $E'_{1\phi}$, $E_{1\phi}$, E'_{1z} and E_{1z} . Eliminating $E'_{1\phi}$ using equation (46), we obtain a first-order differential equation for $E_{1z}(R)$:

$$D_1 E'_{1z} + D_2 E_{1z} + D_3 E_{1R} = 0, \quad (50)$$

where D_1 , D_2 and D_3 are functions of R . The expressions are relatively long and we give them in Appendix B. The $\hat{z}$ component of (49) has no new information; it just gives back equation (44). The $\hat{R}$ component, however, gives a new relation between the various components of $\vec{E}_1$ and their derivatives. Eliminating $E'_{1\phi}$, $E''_{1\phi}$, E'_{1z} and E''_{1z} using equations (46), (47), (50) and the derivative of (50), we obtain a differential equation for $E_{1R}(R)$:

$$D_4 E'_{1R} + D_5 E_{1z} + D_6 E_{1R} = 0, \quad (51)$$

where D_4 , D_5 and D_6 are again functions of R and are given in Appendix B.

We have thus reduced the linear mode analysis problem to a pair of first-order differential equations, (50), (51), for $E_{1z}(R)$ and $E_{1R}(R)$. For convenience, we write down the two equations again:

$$E'_{1z} = -\frac{D_2}{D_1} E_{1z} - \frac{D_3}{D_1} E_{1R}, \quad (52)$$

$$E'_{1R} = -\frac{D_5}{D_4} E_{1z} - \frac{D_6}{D_4} E_{1R}. \quad (53)$$

These equations constitute an eigenvalue problem, where ω is the eigenvalue. By numerically solving the equations with appropriate boundary conditions, we obtain ω for given values of m and k .

The singular points of the equations are located at the radii where $D_1(R)$ and $D_4(R)$ vanish. Anticipating later discussion, we write down here the expression for the quantity $D_1 D_4$:

$$D_1 D_4 = -\frac{g}{\omega R} \left[\frac{(\omega R f - c k R h)^2 + (\omega R g - m c h)^2 - (c k R g - m c f)^2}{(\omega R f - c k R h)} \right]. \quad (54)$$

Also, from equation (10), the perturbed velocity is

$$\frac{\vec{v}_1}{c} = \frac{\vec{E}_1 \times \vec{B}_0 + \vec{E}_0 \times \vec{B}_1}{B^2}, \quad (55)$$

which in component form gives

$$\frac{v_{1R}}{c} = \frac{g E_{1\phi} + f E_{1z}}{f^2 + g^2}, \quad (56)$$

$$\frac{v_{1\phi}}{c} = -\frac{g E_{1R} - h B_{1z}}{f^2 + g^2}, \quad (57)$$

$$\frac{v_{1z}}{c} = -\frac{f E_{1R} + h B_{1\phi}}{f^2 + g^2}. \quad (58)$$

We now consider boundary conditions. A physically valid perturbation will be well-behaved on the axis ($R = 0$) and will satisfy suitable boundary conditions at the jet boundary ($R = R_j$). The condition on the axis is different for axisymmetric ($m = 0$) and non-axisymmetric ($|m| \geq 1$) perturbations, so we consider each of these cases in turn. At the jet boundary, we assume that the jet is constrained by a ‘‘rigid wall’’ and we write down the corresponding boundary condition. In the following, we employ the specific forms of $f(R)$, $g(R)$, $h(R)$ given in equations (16), (17), (18).

3.2. Boundary Condition on the Axis: $m = 0$

Setting $m = 0$ and substituting the expressions for $f(R)$, $g(R)$, $h(R)$ in $D_1 - D_6$, we find that the leading terms of the differential equations (52), (53) at small R are given by

$$E'_{1z} = \frac{a_{zz}}{R} E_{1z} + a_{zR} E_{1R}, \quad (59)$$

$$E'_{1R} = \frac{a_{Rz}}{R^2} E_{1z} + \frac{a_{RR}}{R} E_{1R}, \quad (60)$$

where

$$a_{zz} = -\frac{2\omega}{ck}, \quad (61)$$

$$a_{zR} = \frac{i(c^2 k^2 - \omega^2)}{c^2 k}, \quad (62)$$

$$a_{Rz} = \frac{4i\omega}{k(ck - \omega)}, \quad (63)$$

$$a_{RR} = \frac{(ck + 2\omega)}{ck}. \quad (64)$$

Requiring the perturbation to be analytic as $R \rightarrow 0$ immediately gives the following solution near the axis,

$$E_{1z} = K R^2, \quad (65)$$

$$E_{1R} = -K \frac{2ic}{(ck - \omega)} R, \quad (66)$$

where K is an arbitrary normalization constant.

3.3. Boundary Condition on the Axis: $|m| > 0$

When $m \neq 0$, we obtain

$$a_{zz} = 0, \quad (67)$$

$$a_{zR} = \frac{imA(ck - \omega)}{(Acm - \omega R_m)}, \quad (68)$$

$$a_{Rz} = -\frac{im(Acm - \omega R_m)}{A(ck - \omega)}, \quad (69)$$

$$a_{RR} = -1, \quad (70)$$

where the constant A is defined to be

$$A = [2(\gamma_m^2 - 1)]^{1/2}. \quad (71)$$

The physically relevant solution close to the axis is then

$$E_{1z} = K R^{|m|}, \quad (72)$$

$$E_{1R} = -K \operatorname{sgn}(m) \frac{i(Acm - \omega R_m)}{A(ck - \omega)} R^{|m|-1}. \quad (73)$$

3.4. Boundary Condition at the Jet Boundary: Rigid Wall

We assume that our cylindrical jet is terminated at $R = R_j$ by a rigid impenetrable wall. By impenetrable we mean that no energy flows across this boundary, either out of or into the jet, i.e., the Poynting flux lies in the ϕ - z plane. Equivalently, the velocity vector has no radial component.

The equilibrium Poynting flux of course lies in the ϕ - z plane. The perturbed Poynting flux is proportional to $\vec{E}_1 \times \vec{B}_0 + \vec{E}_0 \times \vec{B}_1$. Since $\vec{E}_0$ is parallel to $\hat{R}$, the term $\vec{E}_0 \times \vec{B}_1$ is automatically in the ϕ - z plane. The term $\vec{E}_1 \times \vec{B}_0$ will also be in this plane if $\vec{E}_1$ is precisely radial, i.e., both $E_{1\phi}$ and E_{1z} vanish. By equation (44), $E_{1\phi}$ is proportional to E_{1z} . We thus obtain the following boundary condition at the outer wall:

$$E_{1z} = 0, \quad R = R_j. \quad (74)$$

4. NUMERICAL RESULTS

We have computed frequencies of modes by numerically solving the differential equations (52) and (53), along with the boundary conditions described in §§3.2–3.4. For each choice of k and m , a countable infinity of solutions exists which may be ordered by the number of zeros of $\text{Re}[E_{1z}(R)]$, not counting zeros at the boundaries.⁹ The lowest-order solution (the “fundamental mode”) is such that $\text{Re}[E_{1z}(R)]$ has no zeros between $R = 0$ and $R = R_j$, the next solution has one zero inside the jet, and so on. In the following we identify each mode by its radial mode number n which we define to be the number of zeros. As one might expect, the mode frequency increases with increasing n .

We solve for the frequencies via a shooting method. We start with a guess value of ω , make use of the expressions given in §3.2 or §3.3 (depending on the value of m) to set up the initial solution at small R , and integrate equations (52) and (53) to $R = R_j$. We then adjust ω in the complex plane until the outer boundary condition given in §3.4 is satisfied. The only subtle point is that the quantities D_1 and D_4 appear in the denominators of various coefficients in equations (52) and (53), and so their zeros correspond to poles in the solution. To avoid these poles, we treat R as a complex variable and integrate the equations over a “safe” trajectory in complex- R space. Since the solution is analytic, the exact track that we follow is unimportant so long as it lies *above* all singularities in the R -plane. Istomin & Pariev (1996) give a detailed discussion of this topic in connection with current-driven instabilities in force-free jets. The reader is also referred to standard discussions of this point in plasma physics texts in the context of Landau damping, or Goldreich, Goodman, & Narayan (1986) for a discussion in the context of accretion disk instabilities.

4.1. Axisymmetric Modes: $m = 0$

Figure 3 shows the dispersion relation — the variation of ω with k — of axisymmetric modes ($m = 0$). Results are shown for Models A, B and C (eqs. 30, 33, 34) for three radial mode numbers: $n = 0, 1, 2$. For all k , we find that the mode frequency is real, which means that all these modes are stable.

For large values of kR_j , the mode frequency asymptotes to $\omega = \pm ck$, so the modes behave like electromagnetic waves moving parallel or anti-parallel to the z -axis. At small k , however, the frequency asymptotes to a constant value. There is thus a minimum frequency for propagating modes inside the jet. The minimum frequency is of order the inverse of the

⁹ $\text{Re}()$ stands for the real part of a complex quantity.




FIG. 3.— Dispersion relation for axisymmetric modes ($m = 0$) in Model A (solid lines), Model B (dashed lines) and Model C (dotted lines). From below the curves correspond to radial mode numbers $n = 0, 1, 2$. All the modes are stable.




FIG. 4.— Imaginary part of ω for modes in Model B with $m = 1, n = 0$. Growing modes have $\text{Im}(\omega) > 0$, while decaying modes have $\text{Im}(\omega) < 0$.

light-crossing time across a radial wavelength of the mode (e.g., $\omega_{\min} \sim 2\pi c/R_j$ for the mode with $n = 0$).

We find that the dispersion relations of modes with positive and negative k are not quite the same. The difference arises because the background has a non-zero velocity in the z direction, which breaks the symmetry between waves propagating towards $+z$ and $-z$. The effect is, however, quite weak.

4.2. Non-Axisymmetric Modes: $m = \pm 1$

The most interesting modes are those with $m = \pm 1$. These modes are stable in Model A, but unstable in Models B and C.




FIG. 5.— Eigenfunctions corresponding to growing modes in Model B with $m = 1$, $n = 0$ and $kR_j = 5, 10, 15, 20$ and 25 . The real part of E_{1z} is plotted.

Figure 4 shows $\text{Im}(\omega)$ ¹⁰ as a function of kR_j for a sequence of modes in Model B; the modes correspond to $m = 1$, $n = 0$. In this sequence, modes with $kR_j < 0.65$ and those with $kR_j > 28$ are stable and have ω real. However, for $0.65 < kR_j < 28$, we find a pair of modes with complex values of ω . The branch with $\text{Im}(\omega) > 0$ corresponds to growing modes, and the branch with $\text{Im}(\omega) < 0$ to decaying modes.¹¹

Figure 5 shows eigenfunctions corresponding to a few of the growing modes. Plotted is $\text{Re}(E_{1z})$ as a function of the scaled radius R/R_j . The mode corresponding to $kR_j = 5$ is representative of all modes with $kR_j \lesssim 5$. These modes have eigenfunctions with no zero crossings between $R = 0$ and $R = R_j$. By our definition, the modes correspond to $n = 0$. Each of the remaining eigenmodes in Fig. 5 has a pronounced dip in $\text{Re}(E_{1z})$ which causes a zero crossing. These dips result from a singularity in the equations, as we discuss below. If we discount the singularity-induced zero crossings, then these eigenfunctions may also be identified as $n = 0$ modes.

Figures 6 and 7 show similar results for Model C. Growing modes (and their decaying counterparts) are present for $m = 1$ and all $kR_j < 2.1 \times 10^4$. Modes with $m = -1$ are also unstable (see Fig. 6)¹². Figure 7 shows a few eigenfunctions. All modes with $kR_j \lesssim 1.3 \times 10^3$ have eigenfunctions with the standard $n = 0$ shape (see the mode with $kR_j = 1.25 \times 10^3$ in Fig. 7). For larger values of k , the eigenfunctions develop negative spikes due to the presence of a singularity (see Fig. 7). However, we still view them as $n = 0$ modes.

We have determined numerically that the singularities which cause the dips in the eigenfunctions are due to zeros in the function $D_4(R)$ defined in §3.1. This function appears in the denominator of the differential equation (53), and hence its zeros behave like poles.¹³

Equation (54) gives the analytic form of the quantity $D_1 D_4$.

¹⁰ $\text{Im}()$ refers to the imaginary part of a complex quantity.

¹¹ We discuss the sudden jump in the value of $\text{Im}(\omega)$ for the decaying branch at the end of §4.2.

¹² In the case of Model B, modes with $m = -1$ appear to be stable, and only the $m = +1$ modes show an instability.

¹³ In contrast, although the function $D_1(R)$ appears in the denominator of equation (52), its zeros do not cause a real singularity since the terms D_2 and




FIG. 6.— Imaginary part of ω for growing modes in Model C. The solid line corresponds to modes with $m = 1$, $n = 0$ and the dotted line corresponds to modes with $m = -1$, $n = 0$.




FIG. 7.— Eigenfunctions corresponding to growing modes in Model C with $m = 1$, $n = 0$ and $kR_j = 1.25 \times 10^3, 2.5 \times 10^3, 5 \times 10^3, 10^4$ and 2×10^4 . The real part of E_{1z} is plotted.

Since the modes of interest to us have $\text{Re}(\omega)$ very nearly equal to ck , let us substitute $\omega = ck$ in this equation. Then, setting $D_1 D_4$ equal to zero gives the following relation between the wavenumber k of the mode and the radius R_{sing} of the singularity:

$$kR_{\text{sing}} = \frac{m[f(R_{\text{sing}}) + h(R_{\text{sing}})]}{g(R_{\text{sing}}) + \text{sgn}(m)\sqrt{g^2(R_{\text{sing}}) + [f^2(R_{\text{sing}}) - h^2(R_{\text{sing}})]}}. \quad (75)$$

Figure 8 shows the position of the singularity R_{sing} as a function of kR_j for modes with $m = \pm 1$ in Models B and C, as calculated with this equation. For comparison, the dots show the radii at which the functions $\text{Re}[E_{1z}(R)]$ reach their minima

D_3 also go to zero at the same locations.




FIG. 8.— Location of the singularity R_{sing} as a function of kR_j in Models B and C, calculated using eq. (75). Solid curves correspond to modes with $m = 1$, $n = 0$ and dotted lines to modes with $m = -1$, $n = 0$. Solid dots show the locations of minima in the eigenfunctions plotted in Figs. 5 and 7.

in the eigenfunctions plotted in Figs. 5 and 7. The agreement between the analytical curve and the dots is excellent, showing that equation (75) captures the physics of the singularity.

From Fig. 8 we see that the singularity lies inside the jet ($R_{\text{sing}} < R_j$) only for a finite range of k above a certain minimum value. For values of k smaller than this minimum, the singularity is outside the jet (for very small k it is well outside the jet). In the case of Model B, the singularity enters the jet from outside when $kR_j \sim 5$ and it disappears (for $m = +1$) at $R = 0$ when $kR_j \sim 28$. This is the primary range of k over which an unstable mode is present. At $kR_j \sim 28$, the singularity is barely present near the center of the jet and we have a very weakly growing mode. With decreasing k , the singularity moves outward and the growth rate of the mode increases (Fig. 4). At $kR_j \sim 5$, when the singularity reaches the wall, the growth rate is close to its maximum value. At yet smaller values of k , the singularity moves outside the outer wall, but its presence is still felt and there is continued instability. The growth rate however decreases with decreasing k .

A similar pattern is seen in Model C. Unstable modes are present only for $kR_j \lesssim 2 \times 10^4$. With decreasing k the growth rate increases and reaches its maximum approximately when the singularity reaches the jet boundary ($R_{\text{sing}} = R_j$), which happens at $kR_j \sim 1.3 \times 10^3$. In contrast to Model B, however, the growth rate remains large even for smaller values of k , and the instability survives down to $k \rightarrow 0$.

We finally discuss the peculiar behavior of $\text{Im}(\omega)$ in the decaying branch of modes in Fig. 4. As we mentioned earlier, in numerically solving for the eigenvalue we must integrate the differential equations (52) and (53) along a path in the complex- R plane that lies above the poles in the solution. For growing modes, the pole is located below the real R -axis. We can therefore integrate along the real R -axis without any difficulty. For decaying modes, the pole is above the real R -axis and now we must choose the integration path with care. If the singularity has $\text{Re}(R_{\text{sing}}) > R_j$, i.e., the singularity is outside the jet, there is no problem and we can simply integrate along

the real axis. However, when $0 < \text{Re}(R_{\text{sing}}) < R_j$, we have to deform the integration path. In our calculations, we integrate from $R = 0$ along a path with $\text{Im}(R) = \text{Re}(R)$ until the point $\text{Im}(R) = \text{Re}(R) = R_j$ and we then integrate down to $R = R_j$. The jump in $\text{Im}(\omega)$ in Fig. 4 is the result of the singularity moving into the jet. To the left of the break, the singularity is located at $R > R_j$. Here the eigenvalues of the growing and decaying modes are complex conjugates of each other. However, to the right of the break, the singularity has moved inside the jet ($R < R_j$) and now the complex conjugate symmetry is broken.

We note that eigenfunctions and eigenvalues of decaying modes are not very meaningful. This can be shown from an initial condition analysis along the lines of Landau's treatment of plasma damping. The reader is referred to Istomin & Pariev (1996) for a detailed discussion of this topic.

4.3. Why $m = \pm 1$ is Special

We have not exhaustively explored modes with $|m| > 1$. However, in spot tests with various choices of m , n and kR_j in Models A, B and C, all modes were found to be stable. We believe that, if at all, there are only weakly unstable modes for $|m| > 1$; there is no sign of the kind of vigorous instability described in §4.2 for modes with $m = \pm 1$. So why is $m = \pm 1$ special? The answer to this question is well-known in the magnetic confinement literature (e.g., Bateman 1978). We discuss it briefly here for completeness.

Consider the radial component of the perturbed velocity v_{1R} near the axis of the jet. Equation (56) gives the expression for v_{1R} in terms of the perturbed electric field components E_{1z} and $E_{1\phi}$, and equation (44) shows the relation between these two field components. For small values of R near the axis, we have

$$g(R) \approx 1, \quad f(R) \approx h(R) = A \frac{R}{R_m}, \quad (76)$$

where the quantity A is defined in equation (71), and

$$C_1 \approx \frac{(\omega R_m - Acm)}{AR(\omega - ck)}. \quad (77)$$

Consider first modes with $m = 0$. Equation (65) shows that $E_{1z} \approx KR^2$ near the axis. Substituting this in equation (56) and using the other approximations given above, we find

$$v_{1R} \approx \frac{\omega R_m}{A(\omega - ck)} KR + \mathcal{O}(R^3). \quad (78)$$

By symmetry, the velocity goes to zero on the axis, and the flow consists of a simple radial divergence.

Consider next modes with $m \neq 0$. Using equation (72) for E_{1z} , we find

$$v_{1R} \approx \frac{(\omega R_m - Acm)}{A(\omega - ck)} KR^{|m|-1} \cos m\phi + \mathcal{O}(R^{|m|+1}), \quad (79)$$

where we have included $\cos m\phi$ to show the angular dependence of the mode. The leading term goes like $R^{|m|-1}$, which corresponds to R^0 when $m = \pm 1$. This means that the mode has a finite radial velocity, and hence a finite radial displacement, on the axis when $|m| = 1$. The $\cos \phi$ dependence of v_{1R} , coupled with the fact that $v_{1\phi}$ has the same amplitude but a $\sin \phi$ dependence, ensures that the velocity vector is unique and analytic at $R = 0$. By writing the velocity vector in cartesian coordinates, it is easily seen that the complex phase of

v_{1R} determines the orientation of the velocity vector in the xy -plane. If we consider values of $|m| \geq 2$, the velocity vanishes on the axis, just as in the case of $m = 0$.

This then reveals what is special about $|m| = 1$ modes. These are the only modes in which fluid perturbations communicate across the axis and cause the jet to shift bodily across the axis. In modes with $|m| = 1$ the center of mass of the jet itself shifts into a spiral shape, which is the characteristic feature of the kink or screw mode. For all other values of $|m|$, the center of mass remains on the axis and the perturbations are concentrated on the outside.

In helical MHD configurations in the laboratory, the $|m| = 1$ kink mode is known to be highly unstable and to be the greatest threat to the stability of equilibria (Bateman 1978). Not surprisingly we see the same feature in our force-free jet equilibria.

4.4. Growth Rate of the Instability

The growth rate of the fastest growing mode is a matter of practical interest since it limits the lifetime of an unstable system. As discussed in §4.2, for the models we have considered here, the most unstable mode generally has a singularity close to the outer wall: $R_{\text{sing}} \sim R_j$. Knowing this, we estimate here the fastest growth rate by assuming that the pole is located at $R_{\text{sing}} = 1.1R_j$. (We locate the singularity slightly outside the jet, since this speeds up the numerical integrations considerably.)

Given an assumed value of R_{sing} , we can substitute this value in equation (75) and make use of the expressions for $f(R)$, $g(R)$, $h(R)$ given in §2.2. Recalling that Models B and C are in the regime described in §2.2.2, we note that $f^2(R_j) - h^2(R_j) \gg g^2(R_j)$. In addition, f and h are nearly equal to each other and γ_j is given by equation (31). We then find that $kR_j \approx 1.6\gamma_j$. Also, the real part of the frequency is nearly equal to ck . Thus, we estimate

$$\text{Mode with } R_{\text{sing}} = 1.1R_j : \quad k \approx \frac{1.6\gamma_j}{R_j}, \quad \text{Re}(\omega) \approx \frac{1.6\gamma_j c}{R_j}. \quad (80)$$

These estimates should apply to the fastest-growing mode. The mode with the maximum growth rate in Model B has $\text{Re}(\omega) \approx kR_j = 4.4$. Since Model B has $\gamma_j = 2.5$, equation (80) predicts $kR_j \approx 4.0$, which is close. Similarly, the mode with the maximum growth rate in Model C has $\text{Re}(\omega) \approx kR_j = 1000$, whereas equation (81) with $\gamma_j = 660$ predicts $kR_j \approx 1060$. We see that the approximate formula (80) is quite good.

Our numerical results indicate that the growth rate $\text{Im}(\omega)$ of the fastest growing mode is proportional to $\text{Re}(\omega)/\gamma_j^2$. We also know that unstable modes are present only when $R_m < R_j$; for instance, Model A with $R_m = 1.2R_j$ has no unstable modes, whereas Model B with $R_m = 0.3R_j$ and Model C with $R_m = 0.18R_j$ both have unstable modes. With these clues in mind, we obtain the following empirical estimate for the growth rate of the fastest-growing mode:

$$\text{Mode with } R_{\text{sing}} = 1.1R_j : \quad \text{Im}(\omega) \approx \frac{0.4}{\gamma_j} \left(1 - \frac{2R_m}{R_j}\right) \frac{c}{R_j}. \quad (81)$$

The coefficients 0.4 and 2 are very approximate (to emphasize this, we give only the leading digit for each). Nevertheless, as we show in Figs. 9 and 10, this approximate formula does quite a good job of fitting the numerical results for a wide range of models. The only region of parameter space where




FIG. 9.— Numerically calculated growth rates of modes with $m = 1$ and $R_{\text{sing}} = 1.1R_j$. These modes have among the largest growth rates. The solid lines show the results for a series of models for fixed γ_m and varying R_m/R_j . The dotted lines are the growth rates predicted by eq. (81). Note the very good agreement except near the top of the plot, where the models are non-relativistic. The dashed lines are the numerical growth rates for modes with $k = 0$.




FIG. 10.— Numerically calculated growth rates of modes with $m = 1$ and $R_{\text{sing}} = 1.1R_j$. The solid lines show the results for a series of models with fixed R_m/R_j and varying γ_m . The dotted lines are the growth rates predicted by eq. (81). The dashed lines are the numerical growth rates for modes with $k = 0$.

the formula fails is when the underlying equilibrium becomes “non-relativistic” and γ_j approaches unity. These models are near the upper end of Figs. 9 and 10 and have extremely large growth rates.

Although modes with $R_{\text{sing}} \sim R_j$ have the largest growth rates, these modes have relatively short wavelengths $\ll R_j$ along the z -axis (see eq. 80). With such short wavelengths it

is not clear if these instabilities can grow to large amplitude. It is therefore interesting to consider modes with $k \rightarrow 0$. Figure 4 shows that Model B is stable as $k \rightarrow 0$, whereas Fig. 6 indicates that the $k = 0$ mode in Model C is nearly as unstable as the fastest-growing mode.

The dashed lines in Figs. 9 and 10 show numerical results for the growth rates of modes with $k = 0$ for various combinations of the model parameters γ_m and R_m . For small values of $R_m \lesssim 0.1R_j$, the results are nearly identical to those we described above for the fastest-growing mode ($R_{\text{sing}} = 1.1R_j$, solid lines). This is to be expected based on the results shown in Fig. 6 for Model C, which has $R_m = 0.1R_j$. With increasing R_m , however, the $k = 0$ modes become less unstable than the modes with $R_{\text{sing}} \sim R_j$. By $R_m \sim 0.3R_j$, the $k = 0$ modes are fully stable, thus explaining the result shown in Model B (Fig. 4), which has $R_m = 0.3R_j$.

4.5. Spatial Growth of Unstable Modes

The discussion so far was limited to modes with real k and complex ω . An equally interesting problem is to consider modes with real ω and complex k . From equation (37), we see that the eigenfunctions take the form

$$\begin{aligned} \vec{E}_1 &= [E_{1R}(R)\hat{R} + E_{1\phi}(R)\hat{\phi} + E_{1z}(R)\hat{z}] \\ &\times \exp[-i\omega t + im\phi + i\text{Re}(k)z - \text{Im}(k)z] \\ &\propto \exp[i\text{Re}(k)z] \exp(z/Z), \end{aligned} \quad (82)$$

where ω is real and $Z = -1/\text{Im}(k)$ is the scale length on which the mode e -folds in the z -direction. Such spatially growing “convected” modes are particularly relevant for sources with long-lived steady jets.

As discussed in Payne & Cohn (1985) and Appl & Camenzind (1992), there is a strong symmetry between modes with real k and complex ω , and those with complex k and real ω . In particular, the growth rates of the two kinds of modes are related by

$$\text{Im}(k) = -\text{Im}(\omega)/v_g, \quad (83)$$

where $v_g = \partial \text{Re}(\omega)/\partial \text{Re}(k)$ is the group velocity of the mode. Our unstable $m = 1$ modes have v_g very nearly equal to c . Therefore, we immediately obtain from equation (81) the following estimate of the spatial e -folding scale Z of the fastest-growing convected mode:

$$\text{Mode with } R_{\text{sing}} = 1.1R_j : \quad \frac{Z}{R_j} \approx \frac{2.5\gamma_j}{(1 - 2R_m/R_j)}. \quad (84)$$

Zero-frequency modes ($\omega = 0$) should have almost the same Z for small values of R_m/R_j , but the growth should cut off at a somewhat smaller value of R_m compared to the modes with $R_{\text{sing}} = 1.1R_j$.

Figure 11 shows numerical results. Modes with $R_{\text{sing}} = 1.1R_j$ have growths consistent with equation (84), and the modes with $\omega = 0$ have similar growths except that the instability cuts off at somewhat smaller values of R_m . The results are as expected and are very similar to those shown in Fig. 9.

The above results correspond to an idealized cylindrical jet. In the case of real jets we must allow for a finite opening angle $\theta_j \equiv dR_j/dz$. Many force-free jet models have opening angles that vary inversely as the Lorentz factor: $\theta_j \sim \text{few}/\gamma_j$ (TMN08). Using this scaling we can estimate approximately the evolution of the mode amplitude a with distance:

$$\begin{aligned} \frac{da}{dR_j} &= \frac{1}{\theta_j} \frac{da}{dz} \approx \frac{\gamma_j}{\text{few}} \frac{da}{dz} \approx \frac{\gamma_j}{\text{few}} \frac{a}{Z} \\ &\approx \frac{\gamma_j}{\text{few}} \frac{a}{2.5\gamma_j R_j} \approx \frac{1}{\text{few} \times 2.5} \frac{a}{R_j}. \end{aligned} \quad (85)$$




FIG. 11.— Numerically calculated e -folding scale Z for modes with $m = 1$, ω real and $R_{\text{sing}} = 1.1R_j$. These modes have among the largest largest growth rates. The solid lines show the results for a series of models with a given value of γ_m and different values of R_m/R_j . The dotted lines are the growths predicted by eq. (84). Note the very good agreement except near the bottom of the plot, where the models are non-relativistic. The dashed lines are the numerical values of Z for modes with $\omega = 0$.

Solving this differential equation, and using $R_j \propto z^{\alpha/4} \sim z^{0.5-1}$ (eq. 6), we obtain

$$a(z) \propto z^\epsilon, \quad (86)$$

where ϵ is a small number $\lesssim 0.1$. This estimate is very crude, but it does suggest that, in realistic jets, the unstable kink mode we have studied in this paper grows only weakly with increasing distance.

4.6. Towards an Improved Instability Criterion

In §1, we introduced three different instability criteria, of which the IPL and TMT criteria refer specifically to rotating force-free jets. Since B_z is practically constant in our equilibria, all of our models are close to the boundary between stability and instability according to the IPL criterion (eq. 2). Similarly, since $B_\phi^2 \approx \Omega^2 R^2 B_p^2$ in our equilibria, our models are marginally stable according to the TMT criterion (eq. 3).¹⁴ It is thus not possible to understand from either of these criteria why Model A is stable and Models B and C are unstable.

It is, of course, not surprising that the above instability criteria fail. Our jet equilibria include the effects of poloidal field curvature, which were not considered by the previous authors. For easier comparison with previous work, let us rewrite our balance condition (22) as follows:

$$\frac{1}{R^2} \frac{d}{dR} \left[\frac{(B_{0\phi}^2 - E_{0R}^2)R^2}{8\pi} \right] = -\frac{d}{dR} \left(\frac{B_{0z}^2}{8\pi} \right) + \left(\frac{E_{0R}^2 - B_{0z}^2}{4\pi R_c} \right). \quad (87)$$

If we leave out the last term, this is equivalent to equation (6) in Istomin & Pariev (1996) and equation (16) in Lyubarskii (1999). The IPL instability criterion states that the quantity

¹⁴ A strict application of the TMT criterion would indicate that our models are unstable, since $B_\phi^2 > E^2 = \Omega^2 R^2 B_p^2$. However, $B_\phi^2 - E^2 \ll B_\phi^2$, so the models deviate only slightly from marginal stability.

dB_{0z}^2/dR should be negative. We might wish to generalize this by saying that the right hand side of equation (87), including the poloidal curvature term, should be positive. Unfortunately, this simple modification is not sufficient since the right hand side is positive for all of our models, whereas not all our models are unstable; not only should the right hand side be positive, its magnitude should be larger than some amount. The same seems to be true with the TMT criterion. This criterion indicates that all our equilibria should be unstable, whereas only some of them are.

Qualitatively, what distinguishes the unstable Models B and C from the stable Model A is that the former have made the transition to the second acceleration regime. This is reflected in their $\gamma(R)$ profiles (Fig. 2) which have $d\gamma/dR < 0$ at larger radii. Thus, one might guess that instability requires the jet to be in the second acceleration regime and/or the jet to have a declining $\gamma(R)$. Once again, these conditions by themselves are not sufficient. To have an instability, $\gamma(R)$ should decline over a sufficiently broad range of radius, e.g., R_m should be less than $\sim 0.45R_j$ in our models.

An alternate approach which we have found useful is to focus on the left-hand side of equation (87). From equations (16), (17), (18), we see that for our equilibria we have $B_{0\phi}^2 - E_{0R}^2 = (R/R_m)^4 B_{0z}^2$. Furthermore, we have seen that modes with $R_{\text{sing}} \sim R_j$ (the fastest-growing modes) are unstable so long as $R_m \lesssim 0.45R_j$, while modes with $k \rightarrow 0$ (long-wavelength modes) are unstable for $R_m \lesssim 0.3R_j$. From this, we obtain the following approximate instability criteria:

$$\text{Modes with } R_{\text{sing}} \sim R_j : (B_{0\phi}^2 - E_{0R}^2)^{1/2} > 5|B_{0z}|, \quad (88)$$

$$\text{Modes with } k \rightarrow 0 : (B_{0\phi}^2 - E_{0R}^2)^{1/2} > 12|B_{0z}|, \quad (89)$$

where we have set $R = R_j$ to obtain the numerical coefficients on the right. These conditions are easier to interpret if we boost to the comoving frame of the fluid (V. Pariev, private communication), where the electric field vanishes. In this frame, the criterion for instability becomes

$$\text{Modes with } R_{\text{sing}} \sim R_j : |B_{0\phi, \text{comov}}| > 5|B_{0z, \text{comov}}|, \quad (90)$$

$$\text{Modes with } k \rightarrow 0 : |B_{0\phi, \text{comov}}| > 12|B_{0z, \text{comov}}|. \quad (91)$$

That is, in the comoving frame, the toroidal field must dominate the poloidal field by more than a certain critical factor.¹⁵ Written in this form, the condition resembles the KS criterion (eq. 1).

5. SUMMARY AND DISCUSSION

The relativistic jet model we have considered in this paper is particularly simple: it is cylindrical, it assumes force-free conditions, and it assumes rigid rotation. Within the limitations of these reasonable approximations, we have attempted to be as close to numerically simulated jets as possible. We include the effect of poloidal field curvature, which is known to play an important role in numerical force-free jets (§2.1), and we choose functional forms for the various field components in the equilibrium (§2.2) to match as closely as possible our previous force-free simulations (TMN08).

Our equilibrium model is described by two parameters: the maximum Lorentz factor γ_m , and the radius at which this maximum is achieved R_m . The ratio of the latter to the jet radius

R_j determines the basic physics of the equilibrium. Models in which $R_m/R_j > 1$ have $\gamma(R)$ increasing monotonically with radius R out to some maximum Lorentz factor $\gamma_j < \gamma_m$ at the outer edge of the jet. Model A (Fig. 2) is an example. In these models the entire jet is in the first acceleration regime (see §2.1, 2.2.1 and TMN08 for details). We find that all these models are perfectly stable.

Models with $R_m/R_j < 1$ are more interesting. Here, $\gamma(R)$ increases upto a maximum value γ_m at $R = R_m$ and then decreases down to a Lorentz factor $\gamma_j < \gamma_m$ at $R = R_j$. Models B and C (Fig. 2) are examples of this kind of model. In these models, the jet fluid at $R < R_m$ is in the first acceleration regime, while the fluid at $R_m < R < R_j$ is in the second acceleration regime. We find that the subset of these models with $R_m/R_j \lesssim 0.45$ are linearly unstable. For R_m/R_j just below 0.45, all the unstable modes have short wavelengths in the z -direction: $\lambda = 2\pi/k_z \sim 2\pi R_j/\gamma_j$. With decreasing R_m , a wider range of k_z becomes unstable, and for $R_m/R_j \lesssim 0.3$, we find that waves with $k_z = 0$, i.e., with arbitrarily long wavelengths, are unstable. The latter modes are perhaps of most interest since they are likely to grow to the largest amplitudes. The numerical results are summarized in Figs. 3–11.

The unstable modes we find are all kink modes with azimuthal wavenumber $m = \pm 1$. These are non-axisymmetric modes in which the jet is distorted helically. A key feature is that, at each z , the center of mass of the jet is shifted away from $R = 0$. It is well-known that MHD configurations with toroidal fields are especially susceptible to the kink mode (Bateman 1978), and our models follow this trend. However, because our equilibria both rotate and move relativistically along z , the criterion for instability is different from the usual KS criterion (eq. 1).

We find that the typical growth rate of the unstable kink mode in our jet models is given by equation (81): the e -folding time is of order γ_j times the light-crossing time R_j/c across the jet. For convected modes with a real frequency, this translates to an e -folding length scale of order γ_j times the jet radius R_j . Since jets typically have opening angles $\sim \text{few}/\gamma_j$, the net result is that the unstable modes grow only slowly with distance from the base of the jet (eq. 86). Of course, relativistic jets in astrophysical sources propagate over many decades, so in principle even this slow growth might lead to a large amplitude of the perturbation. Nevertheless, the fact that the growth is very slow reduces the seriousness of the kink instability.

Our jet equilibria turn out to be close to the boundary between stability and instability according to either the IPL or TMT criterion (eqs. 2, 3), so these criteria are not useful for interpreting the results. In addition, since our models include the effects of poloidal field curvature, they lie outside the range of validity of the IPL and TMT criteria. The most useful instability criterion we have come up with is that, in the comoving frame of the jet fluid, the tangential field should be an order of magnitude or more larger than the poloidal field (eqs. 90, 91). Expressed thus, the criterion is similar to the KS criterion (1), except that it should be applied in the comoving frame and z should be taken to be $\sim R_j$.

All the work described here assumes a rigid wall enclosing the jet at the boundary $R = R_j$. We have done some calculations with a constant pressure boundary and we find unstable modes with much larger growth rates compared to the rigid wall case. However, since we are dealing with a force-free jet, it is not clear that a constant pressure boundary is particularly meaningful. For instance, if the pressure is from a non-

¹⁵ Since our equilibria assume a constant Ω for all field lines, the criteria (90) and (91) are technically valid only for such models. However, since the criteria have been written without any explicit reference to Ω , they may be valid more generally even when Ω varies with R .

relativistic gaseous envelope or cocoon, the gas would have substantial inertia and (we suspect) would probably behave to first approximation like a rigid wall.

Various authors have discussed mechanisms by which instabilities might be suppressed in astrophysical jets. Hardee et al. (2007, and references therein) have shown that an external wind or cocoon can stabilize the Kelvin-Helmholtz mode in MHD jets, though it is not clear if this is relevant for force-free jets. Moll, Spruit, & Obergaulinger (2008) show that lateral expansion causes instabilities to grow more slowly. In a sense, we have already included this effect when we derived the growth rate estimate given in equation (86). In addition, we note that some of the growth suppression seen by Moll et al. is probably because expansion causes different parts of the jet to lose causal contact with one other. This is not an issue for force-free models, where signals propagate at the speed of light.

It would be interesting to simulate numerically the unstable modes described in this paper. Apart from verifying the linear theory, such calculations will reveal the non-linear development of the mode. Does the kink mode saturate at a finite

amplitude and lead to a more-or-less coherent helical pattern or does it destroy the initial equilibrium? This important question can be answered only with 3D simulations. Since the kink mode involves lateral motion of the jet across the axis $R = 0$, the numerical technique used must be flexible enough to allow such motions (e.g., as described by McKinney & Blandford 2008).

We conclude by reminding the reader that the work described here refers to a particularly simple model of relativistic jets which is based on the force-free approximation. In real jets, once the flow crosses the fast magnetosonic point, the inertia of the gas starts to play a role and the force-free approximation is no longer valid (e.g., Tchekhovskoy et al. 2009). In this regime, we must consider the full MHD equations.

The authors thank Alison Farmer for assistance during the early stages of this work and Jonathan McKinney for numerous helpful discussions and comments on the paper. This work was supported in part by NASA grant NNX08AH32G.

APPENDIX

A. CONSTANCY OF POLOIDAL MAGNETIC FIELD ACROSS FORCE-FREE JETS

Figure 2.1 shows that in numerical simulations of force-free jets B_p hardly changes with R . In this Appendix we show that this is a common feature of all jet solutions that smoothly connect to a spinning compact object at the base.

Consider the force balance equation (7). Sufficiently near the compact object, where the jet is in the first acceleration regime (see §2.1), we can drop the terms proportional to R_c^{-1} and $(B_\phi^2 - E^2)$ since in the first acceleration regime $\gamma_1^2 \ll \gamma_2^2$ leading to $E^2/B_p^2 \ll R_c/R$ and $B_\phi^2 - E^2 \ll B_p^2$ (TMN08). Then the force balance equation (7) simplifies to

$$\frac{d(B_p^2)}{dR} \approx 0. \quad (\text{A1})$$

Therefore, sufficiently near the compact object the poloidal field is nearly constant,

$$B_p(R) \approx \text{const}. \quad (\text{A2})$$

Each field line is labeled by the amount of poloidal magnetic flux Φ that it encloses. Due to (A2) this flux can be written simply as

$$\Phi \approx \pi B_p R^2. \quad (\text{A3})$$

These relations are valid throughout the first acceleration regime. We now show that they actually hold asymptotically in all parts of the jet.

Recall that in force-free magnetospheres the enclosed poloidal current I is preserved along each field line (Mestel 1961; Okamoto 1978; Thorne et al. 1986; Beskin 1997; Narayan et al. 2007; Tchekhovskoy et al. 2008),

$$I = I(\Phi) = \frac{c}{2} R B_\phi \approx -\frac{\Omega}{2} B_p R^2, \quad (\text{A4})$$

where the approximate equality is due to $B_\phi \approx -E = -\Omega R B_p / c$ for $R \gg R_A$ (see eq. 11). Comparing (A3) and (A4) and recalling that Ω is conserved along field lines, we obtain

$$\Phi \approx -\frac{2\pi}{\Omega} I(\Phi). \quad (\text{A5})$$

Since both sides of this relation depend only on Φ , this relation is valid *everywhere* in the solution, even though we derived it only in the first acceleration regime. Using (A4) to substitute for I yields back (A3). Thus eqs. (A2) and (A3) are valid everywhere in the jet.

B. COEFFICIENTS IN EQUATIONS (52) AND (53)

In this Appendix we give explicit expressions for the coefficients $D_1(R)$ – $D_6(R)$ defined in §3.1. The functions are

$$D_1(R) = -\frac{ck}{\omega} g - \frac{cm}{\omega R} g C_1, \quad (\text{B1})$$

$$D_2(R) = -\frac{cm}{\omega R^2} f - \frac{cm}{\omega R} f' + C_2 C_1 - \frac{cm}{\omega R} g C_1', \quad (\text{B2})$$

$$D_3(R) = \left(\frac{ick^2}{\omega} + \frac{icm^2}{\omega R^2} - \frac{i\omega}{c} \right) g, \quad (\text{B3})$$

$$D_4(R) = \frac{ck}{\omega} f + \frac{cm}{\omega R} g - h - C_3 \frac{D_3}{D_1}, \quad (B4)$$

$$D_5(R) = C_4 + C_3 \frac{D_2^2}{D_1^2} + \left(C_6 + \frac{C_5}{R} \right) C_1 + C_5 C_1' - C_7 \frac{D_2}{D_1} + C_3 \left(\frac{D_2 D_1'}{D_1^2} - \frac{D_2'}{D_1} \right) + \frac{ic}{\omega} g C_1'', \quad (B5)$$

$$D_6(R) = C_8 + C_3 \frac{D_2 D_3}{D_1^2} - C_7 \frac{D_3}{D_1} + C_3 \left(\frac{D_3 D_1'}{D_1^2} - \frac{D_3'}{D_1} \right), \quad (B6)$$

where primes denote derivatives with respect to R , and the functions $C_1(R)$ – $C_8(R)$ are given by

$$C_1(R) = \frac{\omega R g - cm h}{\omega R f - ck R h}. \quad (B7)$$

$$C_2(R) = \frac{ck}{\omega R} f - \frac{cm}{\omega R^2} g - \frac{h}{R} - \frac{h}{R_c} + \frac{ck}{\omega} f' - h', \quad (B8)$$

$$C_3(R) = \frac{ic}{\omega} (f + g C_1), \quad (B9)$$

$$C_4(R) = -\frac{icm^2}{\omega R^2} f + \frac{i\omega}{c} f + \frac{ickm}{\omega R} g - ikh, \quad (B10)$$

$$C_5(R) = \frac{ic}{\omega R} g + \frac{ic}{\omega R_c} g + \frac{ic}{\omega} g', \quad (B11)$$

$$C_6(R) = \frac{ickm}{\omega R} f - \frac{ick^2}{\omega} g - \frac{2ic}{\omega R^2} g + \frac{i\omega}{c} g - \frac{im}{R} h, \quad (B12)$$

$$C_7(R) = \frac{2ic}{\omega R} f + \frac{ic}{\omega} f' + C_5 C_1 + \frac{2ic}{\omega} g C_1', \quad (B13)$$

$$C_8(R) = \frac{2ck}{\omega R} f - \frac{cm}{\omega R^2} g - \frac{2h}{R} + \frac{cm}{\omega R_c R} g - \frac{h}{R_c} + \frac{ck}{\omega} f' + \frac{cm}{\omega R} g' - h'. \quad (B14)$$

REFERENCES

- Appl, S. & Camenzind, M. 1992, A&A, 256, 354
 Appl, S., Lery, T., & Baty, H. 2000, A&A, 355, 818
 Bateman, G. 1978, MHD instabilities (Cambridge, Mass., MIT Press, 1978. 270 p.)
 Begelman, M. C. 1998, ApJ, 493, 291
 Benford, G. 1981, ApJ, 247, 792
 Beskin, V. S. 1997, Soviet Physics Uspekhi, 40, 659
 Ferrari, A., Trussoni, E., & Zaninetti, L. 1978, A&A, 64, 43
 Gammie, C. F., McKinney, J. C., & Tóth, G. 2003, ApJ, 589, 444
 Goldreich, P., Goodman, J., & Narayan, R. 1986, MNRAS, 221, 339
 Goldreich, P. & Julian, W. H. 1969, ApJ, 157, 869
 Goldreich, P. & Tremaine, S. 1978, ApJ, 222, 850
 Hardee, P., Mizuno, Y., & Nishikawa, K.-I. 2007, Ap&SS, 311, 281
 Istomin, Y. N. & Pariev, V. I. 1996, MNRAS, 281, 1
 Kadomtsev, B. B. 1966, Reviews of Plasma Physics, 2, 153
 Lery, T., Baty, H., & Appl, S. 2000, A&A, 355, 1201
 Li, L.-X. 2000, ApJ, 531, L111
 Lynden-Bell, D. 2006, MNRAS, 369, 1167
 Lyubarskii, Y. E. 1999, MNRAS, 308, 1006
 McKinney, J. C. 2006, MNRAS, 368, 1561
 McKinney, J. C. & Blandford, R. D. 2008, preprint (arXiv:0812.1060)
 McKinney, J. C. & Gammie, C. F. 2004, ApJ, 611, 977
 Mestel, L. 1961, MNRAS, 122, 473
 Mignone, A. & McKinney, J. C. 2007, MNRAS, 378, 1118
 Mizuno, Y., Hardee, P., & Nishikawa, K.-I. 2007, ApJ, 662, 835
 Moll, R., Spruit, H. C., & Obergaulinger, M. 2008, A&A, 492, 621
 Narayan, R., Goldreich, P., & Goodman, J. 1987, MNRAS, 228, 1
 Narayan, R., McKinney, J. C., & Farmer, A. J. 2007, MNRAS, 375, 548
 Okamoto, I. 1978, MNRAS, 185, 69
 Payne, D. G. & Cohn, H. 1985, ApJ, 291, 655
 Ruderman, M. A. & Sutherland, P. G. 1975, ApJ, 196, 51
 Tchekhovskoy, A., McKinney, J. C., & Narayan, R. 2007, MNRAS, 379, 469
 —. 2008, MNRAS, 388, 551 (TMN08)
 —. 2009, ApJ, submitted (arXiv:0901.4776)
 Thorne, K. S., Price, R. H., & MacDonald, D. A. 1986, Black holes: The membrane paradigm (Black Holes: The Membrane Paradigm)
 Tomimatsu, A., Matsuoka, T., & Takahashi, M. 2001, Phys. Rev. D, 64, 123003